arXiv:1210.0745v1 [quant-ph] 2 Oct 2012

Classical simulation of partially entangled states

H. M. Bharath[1]* and V. Ravishankar[1,2]†
[1] *Department of Physics, Indian Institite of Technology, Kanpur - 208016 (India) and*
[2] *Department of Physics, Indian Institute of Technology, Delhi -110016 (India)*

We investigate the possibility of simulating partially entangled two qubit states by separable states of higher spins. First, we show that all partially entangled isotropic states can be simulated classically. We further investigate partially entangled pure states, and their binary mixtures and find that these systems forbid such a simulation, signifying that separability does not have a universal character in determining the inherent quantum nature of a state.

Introduction: The problem of distinguishing the truly quantum properties of a system from those which can be realized in classical systems has received close attention ever since the birth of quantum mechanics. In particular, the violation of Bell inequality by fully entangled states establishes that there is no classical, i.e., a separable bipartite state that can simulate the spin correlations fully. This raises the interesting question whether there are classical systems which can simulate quantum systems which are partially entangled. The question gains further importance in quantum computation, as may be seen from the remarkable work of Valiant [1] who showed how a class of quantum computers can be simulated by classical computers efficiently – in polynomial time. With these in mind, in this letter, we examine whether partially entangled two qubit states admit a classical simulation. To make the results transparent, and keep the arguments unclutterd, we focus on three classes of states – which suffice to bring out the rich interplay between entanglement and classical quantum correspondence.

Motivation and formulation: Let us start with a quick recap of the concepts of of separability [2] and spin coherent states [3] (which we employ for formulating the problem). A bipartite state is separable if it admits a resolution

$$\rho^{AB} = \sum_i p_i \rho_i^A \otimes \rho_i^B. \tag{1}$$

In such a state, the expectation value of an observable $O_{12} \equiv O_1(A)O_2(B)$ has the property

$$Tr\{O_{12}\rho^{AB}\} = \sum_i p_i Tr\{O_1(A)\rho_i^A\} Tr\{O_2(B)\rho_i^B\} \tag{2}$$

which has precisely the characterestics of a classical (local hidden variable) correlation. States which do not admit such an expansion are nonseparable. The essence of this definition is that any classical bivariate probability density $p(\alpha,\beta)$ is separable; it can be expanded as $p(\alpha,\beta) = \sum_i \lambda_i f_i(\alpha) g_i(\beta)$ in terms of univariate probability densities f_i, g_i and appropriate weights λ_i. The notion of nonseparability implies that there are probability amplitudes which do not share this property.

To make further progress, it is convenient to employ spin coherent basis (SCB). Recall again that spin coherent states $|\hat{n}\rangle$ for a spin S particle may be generated by the action of the rotation group (more precisely, $SU(2)$) on the state $|m = S\rangle$ [3], as shown below:

$$|\hat{n}(\theta,\phi)\rangle \equiv e^{-iS_z\phi} e^{-iS_y\theta} e^{-iS_z\psi} |S\rangle \tag{3}$$

Note that we have ignored the over all phase introduced by $e^{-iS_z\psi}$. The coherent states satisfy the relations $\langle \hat{n}(\theta,\phi)|\vec{S}|\hat{n}(\theta,\phi)\rangle = S\hat{n}(\theta,\phi)$ which justifiies the nomenclature. The set $\{\hat{n}(\theta,\phi)\}$, which constitues SCB, is overcomplete. Yet, the inner product $|\langle \hat{n}|\hat{n'}\rangle| = (\frac{1+\hat{n}\cdot\hat{n'}}{2})^S \rightarrow 0$ as $S \rightarrow \infty$, i.e., the states become mutually orthogonal. Since the manifold of coherent states is simply the phase space for a classical spin, one says that the classical limit is attained in the large S limit.

Central to our analysis is the result that given a state of a given spin S, it is possible to characterize it completely in terms of its diagonal elements in SCB, $F(\hat{n}) = \langle \hat{n}|\rho|\hat{n}\rangle$, meaning that the parent state is uniquely determined by the classical *probability density function* (PDF), $F(\hat{n})$. Thus, for example, a single qubit state has a unique PDF in the manner shown below:

$$\rho = \frac{1}{2}(1 + \vec{\sigma}\cdot\vec{P}) \leftrightarrow F(\hat{n}) = \frac{1}{4\pi}(1 + \vec{P}\cdot\hat{n}). \tag{4}$$

This result extends to a two qubit state ρ^{AB}; its corrresponding PDF is given by

$$F(\hat{n}_A,\hat{n}_B) = \langle \hat{n}_A \otimes \hat{n}_B|\rho^{AB}|\hat{n}_A \otimes \hat{n}_B\rangle \tag{5}$$

and, once again, $F(\hat{n}_A,\hat{n}_B)$ determines the parent state unambiguously.

The stage is set to formulate the probem. Being a classical Liouville function, $F(\hat{n}_A,\hat{n}_B)$ is separable; it admits a resolution

$$F(\hat{n}_A,\hat{n}_B) = \sum_i p_i f_i(\hat{n}_A) g_i(\hat{n}_B) \tag{6}$$

The density functions $f_i(\hat{n}^A)$ and $g_i(\hat{n}^B)$ correspond to some parent states ρ_i^A and ρ_i^B. This would appear to yield a separable resolution of ρ^{AB}

$$\rho^{AB} = \sum_i p_i \rho_i^A \otimes \rho_i^B \tag{7}$$

---

*Electronic address: bharathm@iitk.ac.in
†Electronic address: vravi@iitk.ac.in, vravi@physics.iitd.ac.in

even if the state is nonseparable, leading to an apparent contradiction, as depicted in Fig.1.



Figure 1: Schematic diagram indicating the apparent contradiction.

As mentioned, we illustrate the resolution of this apparent contradiction for three representative cases in the following.

1. Isotropic states: Isotropic two qubit states (Werner states) have the form

$$\rho^W[\alpha] = \frac{1}{4}\{\mathbf{1} \otimes \mathbf{1} + \alpha \vec{\sigma}^A \cdot \vec{\sigma}^B\} \tag{8}$$

where, the parameter $\alpha \in [-1, \frac{1}{3}]$. The corresponding PDF is given by

$$\langle \hat{n}_A \otimes \hat{n}_B | \rho^W[\alpha] | \hat{n}_A \otimes \hat{n}_B \rangle = \frac{1}{4}\{1 + \alpha \hat{n}_A \cdot \hat{n}_B\} \equiv F^W \tag{9}$$

It is important to note that F^W is a legitimate PDF over a larger range of the parameter, i.e., $\alpha \in [-1, 1]$. Being classical, it is separable everywhere. The parent state is, however, separable only when $\alpha \in [-\frac{1}{3}, \frac{1}{3}]$, as follows from the PPT criterion [4, 5].

We show that the separability of F^W in Eq. 8 over the full range of values corresponds, in fact, to the separability of higher spin states, which yield exactly the same correlations as the original two qubit state. It suffices to consider the family of bipartite states where A has spin half and B possesses a spin S, for which we establish the range of separability. Consider the class of isotropic states

$$\rho^{\frac{1}{2},S}[\alpha] = \frac{1}{2(2S+1)}\{\mathbf{1}_{\frac{1}{2}} \otimes \mathbf{1}_S + \alpha \vec{\sigma} \cdot \vec{T}\} \tag{10}$$

where, $\vec{T} = \frac{\vec{S}}{S}$. Positivity constraints imply that $\alpha \in [-1, \frac{S}{S+1}]$, which contains the range for the Werner states as its subset. Furthermore, the PDF for $\rho^{\frac{1}{2},S}[\alpha]$ is again given by F^W but over the larger range, indicating the resolution of the paradox. That this is the case follows from the following result:

Result : The isotropic states, $\rho^{\frac{1}{2},S}[\alpha]$ defined in Eq.9 are separable in the range $|\alpha| \leq \frac{S}{S+1}$[7].

Proof: The proof is constructive, and exploits the isotropy of the states. Consider the pure product state $\rho \equiv (|\frac{1}{2}\rangle \otimes |S\rangle)(\langle\frac{1}{2}| \otimes \langle S|)$. Written in the irreducible tensor basis, this has the form

$$\rho_z = \frac{1}{2}\{\mathbf{1}_{\frac{1}{2}} + \sigma_z\} \otimes \frac{1}{2S+1}\{\mathbf{1}_S + \frac{3S}{S+1}T_z + \sum_{k=2}^{2S} q_z^k T_z^k\} \tag{11}$$

which emphasizes that the quantization is along the z axis. Note that the spin S state admits polarizations of all ranks. From ρ_z, we construct the isotropic state which is done in two steps: (i) We first construct the separable state $\rho^1 = \frac{1}{3}(\rho_x + \rho_y + \rho_z)$ obtained by summing over the three orthogonal directions. This has the form

$$\bar{\rho} = \frac{1}{4S+2}\{\mathbf{1}_{\frac{1}{2}} \otimes \mathbf{1}_S + \frac{S}{S+1}\vec{\sigma} \cdot \vec{T}\} + \cdots \tag{12}$$

where the anisotropic terms, all of higher rank, are indicated by ellipses. (ii) The anisotropic correlation terms in $\bar{\rho}$ are eliminated by a uniformization procedure, i.e., by averaging over all possible rotations, which of course leave the isotropic terms unchanged, and leave us with $\rho^{\frac{1}{2},S}[\alpha]$ [8]. Thus the state $\rho^{\frac{1}{2},S}[\alpha]$ is separable over the range $\alpha \geq -\frac{S}{S+1}$ for any spin S. Since PPT criterion is necessary, this state is nonseparable over the range $\alpha \leq -\frac{S}{S+1}$ [9].

The entangling segment shrinks to increasingly smaller intervals $\alpha \in [-1, -\frac{S}{S+1})$ as S increases. Thus all isotropic states, with the exception of the extremal Bell state can be simulated by a classical state of sufficiently large spin, whose minimum value is given by $S_{min} = -\alpha/(1+\alpha)$. It is understood that if S_{min} lies between two half integers,, the larger half integer is to be chosen. Thus the entanglement of Werner states, as represented by the corresponding S_{min}, brings out the nonclassicality vividly, with $S_{min} \to \infty$ rapidly as $\alpha \to -1$. Fig 2 shows this function in the range $\alpha \in [-0.85, -0.95]$.



Figure 2: The higher spin classical equivalents of Werner states as a function of α.

2. Pure states: The discussion of the isotropic states raises the possibility of partially entangled pure states also admitting a classical simulation. For, none of the partially entangled isotropic states is pure. If that were to be so, entanglement (in the sense of nonseparabilty) would have the character of a universality. The analysis below dispels such a hope. In anticipation of this result, we shall call all those states which do not admit a classical simulation, exceptional. The result is that all partially entangled states are exceptional. The proof involves demonstrating that every bipartite state of two equal spins S, which is equivalent an an entangled two qubit state, is nonseparable; more explicitly, it fails to satisfy the PPT criterion.

All pure states of a given entanglement are related by local $SU(2) \times SU(2)$ transformations. Hence it is convenient to employ the Schmidt basis and write $|\psi(\theta)\rangle = \cos\theta|\uparrow\uparrow\rangle + \sin\theta|\downarrow\downarrow\rangle$. Its corresponding PDF is easily seen to be

$$\begin{aligned} F_\theta(\hat{n}^A, \hat{n}^B) &= \{1 + \cos 2\theta(n_z^A + n_z^B) + n_z^A n_z^B \\ &\quad + \sin 2\theta(n_x^A n_x^B - n_y^A n_y^B)\} \end{aligned} \tag{13}$$

The equivalent higher spin bipartite state is

$$\begin{aligned} \rho &= \frac{1}{(2S+1)^2}\{\mathbf{1}_S \otimes \mathbf{1}_S + \cos 2\theta(T_0^1 \otimes \mathbf{1}_S + \mathbf{1}_S \otimes T_0^1) \\ &\quad + T_0^1 \otimes T_0^1 + \sin 2\theta(T_+^1 \otimes T_+^1 + T_-^1 \otimes T_-^1)\} \end{aligned} \tag{14}$$

where $T_0^1 = S_z/S$; $T_\pm^1 = S_\pm/(\sqrt{2}S)$. Its partially transposed form is given by

$$\begin{aligned} \rho' &= \frac{1}{(2S+1)^2}\{\mathbf{1}_S \otimes \mathbf{1}_S + \cos 2\theta(T_0^1 \otimes \mathbf{1}_S + \mathbf{1}_S \otimes T_0^1) \\ &\quad + T_0^1 \otimes T_0^1 + \sin 2\theta(T_+^1 \otimes T_-^1 + T_-^1 \otimes T_+^1)\}. \end{aligned} \tag{15}$$

If ρ' were positive, so would $\Pi\rho'\Pi$ be, where Π is any projection operator. On the other, the projection in the subspace $\mathcal{H}_2$ spanned by $\{|S,-S\rangle,\ |S-1,-S+1\rangle\}$, has the form

$$\rho_2'(\theta) = \begin{pmatrix} 0 & \frac{Sin2\theta}{S} \\ \frac{Sin2\theta}{S} & 1 - \frac{2S+1}{S^2} \end{pmatrix} \tag{16}$$

which is clearly nonpositive, unless $\sin 2\theta = 0$, in which case, it is trivially separable, at every spin including $S = 1/2$.

3. A class of rank two states: We have seen that all partially entangled isotropic states are mixed and may be simulated classically. It is also settled that partially entangled pure states cannot be simulated classically. To show that the interplay of entanglement and mixedness is more involved, we construct a class of rank two mixed states which are again exceptional. Consider an incoherent mixture of two pure states, $|\psi(\theta)\rangle$ and $|\psi(\theta')\rangle$ in the same Schmidt basis, with respective weights λ and $1-\lambda$. The associated bipartite state of two spin S particles is constructed exactly the same way. The projected state ρ^P in the two dimensional subspace spanned by $|S,-S\rangle$ and $|S-1,-S+1\rangle$ has the form $\rho_2' = \lambda\rho_2'(\theta) + (1-\lambda)\rho_2'(\theta')$, which again, is nonpositive almost everywhere. Thus, all pure entangled states, and some rank two entangled states are genuinely exceptional, emphasizing the importance of coherence alongside entanglement.

Conclusion: We have demonstrated how an isotropic state can, in fact, be simulated classically unless it is a Bell state. The entanglement is directly related to the spin of the separable state, which diverges as the entanglement approaches its maximum. However, a naïve generalization of this result to all partially entangled states would be in error, as we showed through the examples of pure states and their rank two incoherent mixtures. These results emphasize that there is a nontrivial interplay of entanglement and coherence in determining the "quantumness" of a system, which merits further study. Indeed,the field of classical quantum correspondence, and its relation to nonlocality as initiated by Bell[6], remains a fertile ground for further insights.

---

[1] L. Valiant, *in Proceedings of the ACM Symposium on the Theory of Computing* ACM Press, New York, 2001!, p. 114; see http://www.deas.harvard.edu/ valiant. For a physical realization in terms of noninteracting fermions, see Barbara. M. Terhal and David. P. Divincenzo, *Phys. Rev. A* 65, 032325 (2002)

[2] R. F. Werner, *Phys. Rev. A* 40, 4277 (1989)

[3] A. Perelomov, *Generalized coherent states and their applications*, (Springer Verlag, Berlin, 1986) Chapter 4, p54

[4] A. Peres, *Phys. Rev. Lett.* 77:1413 (1996)

[5] M. Horodecki, P. Horodecki, R. Horodecki, *Physics Letters A* 223, 1 (1996)

[6] J S Bell, *Physics* 1, 195 (1964)

[7] Note that $\rho^{\frac{1}{2},S}[\frac{S}{S+1}]$ is essentially a projector onto the spin $(S+\frac{1}{2})$ space.

[8] In practice, one needs only an averaging over a finite number of terms which depend on S, but that is not our prime concern here.

[9] Interestingly in this case, the PPT criterion has proved to be sufficient as well.